

altenpflege.de Edition

Short Textbooks for Nursing Education - Focus on Elderly Care

Theo Ahrens

Pressure Ulcer Prevention and Treatment

presented by

YES
ARCH
ITEC
TURE.

Bibliographic Information

Theo Ahrens – Short Biography

Theo Ahrens, born in 1979 near Munich, is a teacher in nursing professions and a dedicated nursing educator. After completing a voluntary social year in a hospital, he trained as a health and nursing professional, gaining valuable professional experience in the intensive care unit. Driven by his desire to impart knowledge, he studied nursing education and began teaching at a vocational school in Munich in 2009. Theo is known for his practical and empathetic teaching, preparing future nurses for their demanding profession. Privately, he lives with his wife and two children in Berlin and volunteers for better conditions in nursing education.

This short textbook was formulated from the lesson plans of his classes using A.I. The translation was done automatically.

Published by altenpflege.de

Price DE/AT: €3.90 CH: 5 Fr

Copyright 2024 altenpflege.de

Please send feedback and suggestions for improvement to team@altenpflege.de.

1/5-2412181531

preview

Table of Contents

1 Introduction 4	
1.1 Objectives of this Short Textbook 4	
1.2 Importance of Pressure Ulcer Prevention and Treatment in Elderly Care 4	
1.3 Role of Nursing Staff in the Prevention and Treatment of Pressure Ulcers 5	
2 Fundamentals of Pressure Ulcers 6	
2.1 What is Pressure Ulcer 6	
2.2 Pathophysiology of Pressure Ulcers 6	
2.3 Risk Factors for the Development of Pressure Ulcers 7	
2.4 Classification of Pressure Ulcer Stages 7	
2.4.1 Stage 1: Reddening of the skin 7	
2.4.2 Stage 2: Skin Defect 8	
2.4.3 Stage 3: Deeper Tissue Damage 9	
2.4.4 Stage 4: Deep Tissue Injury 10	
3 Causes and Risk Factors for Pressure Ulcers 11	
3.1 Primary Causes 11	
3.2 Secondary Causes 11	
3.3 Individual Risk Factors 12	
3.4 Systemic Factors 12	
4 Diagnosis and Identification of Pressure Ulcers 14	
4.1 Observation and Identification of Early Signs 14	
4.2 Distinguishing Skin Redness 14	
4.3 Use of Assessment Tools 15	
4.4 Documentation of Pressure Ulcers 15	
5 Prophylaxis of Pressure Ulcers – Preventive Measures 17	
5.1 Fundamentals of Pressure Ulcer Prevention 17	
5.2 Risk Assessment and Prevention Planning 17	
5.3 Positioning and Repositioning of Residents 18	
5.3.1 Soft Positioning and Micro-positioning 18	
5.3.2 30° positioning, 135° positioning, free positioning 19	
5.3.3 Mobilization and Activation of Residents 19	
5.4 Pressure Relief through Aids 20	
5.5 Promoting Self-Mobilization and Active Movement Encouragement 21	
5.6 Skin Care and Skin Protection 22	
5.7 Nutrition for Pressure Ulcer Prevention 22	
6 Aids for Pressure Ulcer Prevention 24	
6.1 Positioning Aids 24	
6.2 Pressure Relief Systems 24	
6.3 Seat Cushions and Wheelchair Cushions for Pressure Ulcer Prevention 25	
6.4 Assistive Devices for Mobilization 25	
6.5 Protective Aids for Vulnerable Body Areas 26	
7 Treatment of Pressure Ulcers – Measures and Nursing Interventions 27	
7.1 Care of Pressure Ulcers in Stages 1 to 4 27	
7.2 Wound Cleaning 28	
7.3 Modern Wound Care 28	
7.4 Treatment of Infections in Pressure Ulcers 29	
7.5 Measures to Promote Wound Healing 30	
7.6 Use of Special Care Products 31	
8 Care and Support of Residents with Pressure Ulcers 32	
8.1 Nursing Interventions for Residents with Pressure Ulcers 32	
8.2 Promoting Active Participation of Residents in Care 33	
8.3 Communication with Relatives 33	
8.4 Psycho-Social Care for Residents with Pressure Ulcers 34	
9 Documentation of Prophylaxis and Treatment 36	
9.1 Importance of Documentation 36	
9.2 Contents of Pressure Ulcer Documentation 36	
9.3 Use of Documentation Software for Care Planning 37	
9.4 Photo Documentation of Pressure Ulcers 38	
10 Roles and Responsibilities of Caregivers 39	
10.1 Responsibilities of Caregivers in the Area of Pressure Ulcer Prevention 39	
10.2 Involvement of Caregivers in Risk Assessment and Evaluation 40	
10.3 Interdisciplinary Collaboration with Doctors, Wound Managers, and Therapists 40	
10.4 Importance of Continuous Education for Nursing Staff in Pressure Ulcer Prevention 41	
11 Ethical and Legal Aspects of Pressure Ulcer Prevention 43	
11.1 Right of Self-Determination of Residents 43	
11.2 Liability Issues in Pressure Ulcers 44	
11.3 Responsibility of the Care Facility 44	
11.4 Obligation to Adhere to Pressure Ulcer Prophylaxis According to SGB XI 45	
12 Quality Assurance and Hygiene Measures in Pressure Ulcer Care 47	
12.1 Importance of Quality Assurance in Pressure Ulcer Prevention 47	
12.2 Guidelines on Hygiene and Aseptic Wound Care 47	
12.3 Monitoring and Controlling the Effectiveness of Preventive Measures 48	
12.4 External and Internal Audits for Reviewing Pressure Ulcer Prevention 49	
13 Case Studies and Practical Applications 51	
13.1 Case Study: Care Planning for Pressure Ulcer Prevention in Immobile Residents 51	
13.2 Case Study: Care of a Pressure Ulcer (Stage 2) with Modern Wound Management 52	
13.3 Reflection and Discussion of Nursing Practice in the Treatment of Pressure Ulcers 53	

1 Introduction

1.1 Objectives of this Short Textbook

This concise textbook imparts comprehensive knowledge on pressure ulcer prevention and treatment. It facilitates an understanding of the development, risk factors, and classification of pressure ulcers. The focus is on preventive measures and the professional care of affected individuals. It also outlines effective wound care strategies and the use of aids.

It specifically enhances practical skills through case studies and practical instructions. It enables caregivers to make evidence-based decisions and create individualized care plans. Particular emphasis is placed on interdisciplinary collaboration.

The textbook takes into account the ethical and legal aspects of care. It also includes guidelines for documenting measures—essential for quality assurance. The goal is to support caregivers in the optimal care of residents and sustainably reduce pressure ulcer rates.

1.2 Importance of Pressure Ulcer Prevention and Treatment in Elderly Care

Decubitus ulcers present a significant health risk in elderly care. If left untreated, they can lead to severe health consequences, including infections and chronic wounds. Therefore, prevention requires particular attention.

Decubitus ulcer prevention not only reduces health complications but also promotes the residents' quality of life. Effective pressure relief and diligent skin care are central to minimizing the risk of developing pressure sores. Preventive measures help decrease the care needs and, in the long run, reduce care costs.

In addition to prevention, the early detection and treatment of decubitus ulcers are crucial. Treatment methods such as modern wound care significantly contribute to healing and pain reduction. Multidisciplinary approaches, including collaboration with medical professionals, improve treatment outcomes.

The role of caregivers here is of special importance. Their trained eye for identifying risk factors and the application of professional care practices are essential for successful prevention and treatment. Schools and educational institutions for nursing professions provide guidance to expand these competencies, offering residents the best possible care. Through continuous qualification, caregivers remain up-to-date with the latest scientific knowledge, which is crucial for the successful implementation of preventive and treatment measures.

1.3 Role of Nursing Staff in the Prevention and Treatment of Pressure Ulcers

Caregivers play a central role in the prevention and treatment of pressure ulcers. Their influence begins with the assessment and identification of risk factors. They conduct regular skin inspections and use assessment tools to identify at-risk residents.

Careful planning is crucial. Caregivers create individualized care plans and integrate all necessary measures to minimize risk. The selection of appropriate positioning techniques and intervals is tailored to the resident's specific health condition.

In practice, the use of aids is essential. Caregivers use special mattresses or cushions to reduce pressure on vulnerable areas of the body. The caregiver's involvement is indispensable for properly implementing and adjusting preventive measures.

Mobility is an essential element of prophylaxis. Caregivers actively promote residents' self-mobilization, regularly assist with movement exercises, and integrate mobilization strategies into daily care routines.

In the treatment of existing pressure ulcers, caregivers employ qualified wound care methods. They select suitable dressing materials and care products, ensure hygienic conditions, and serve as key contacts in both direct wound care and communication with doctors and therapists.

The training and continuing education of caregivers are crucial to keep their expertise current. This is the only way they can work effectively in an interdisciplinary team and continuously improve resident care.

Finally, caregivers are responsible for comprehensive documentation. They record changes in skin condition and document all prophylactic and therapeutic measures. This not only ensures the quality of care but also provides the basis for reviewing and adjusting care planning.

2 Fundamentals of Pressure Ulcers

2.1 What is Pressure Ulcer

Decubitus, also known as pressure ulcer, is a localized injury to the skin and underlying tissue caused by prolonged pressure or pressure combined with shear forces. These tissue damages commonly occur in areas where bones are close to the skin surface, such as the heels, sacrum, elbows, or shoulder blades. Due to the pressure, blood circulation in the affected areas is interrupted, leading to inadequate supply of oxygen and nutrients to the tissue. This undernourishment can ultimately lead to tissue death, also known as necrosis. Decubitus is a serious complication that particularly occurs in immobile or bedridden individuals. Early detection and consistent prevention are crucial to avoiding the development of pressure ulcers and ensuring the quality of life for those affected.

2.2 Pathophysiology of Pressure Ulcers

A pressure ulcer primarily develops due to prolonged pressure on the skin and the underlying tissue. This pressure leads to a disruption of the microcirculation. The pressure compresses the capillary vessels, which reduces blood flow and impairs the supply of oxygen and nutrients to the tissue. There is an accumulation of metabolic waste products, which further disrupts cellular metabolism.

Irreversible tissue damage can occur after just two hours of uninterrupted pressure. However, the severe consequences are not solely due to pressure. Other factors such as shear forces, caused by the sliding of skin layers against each other, promote the injury to capillaries, further impair blood flow, and thus contribute to the development of a pressure ulcer.

Vasoactive cytokines and inflammatory mediators are released in response to cell damage. These enhance the local inflammatory reaction. With continued pressure and without intervention, the damage progresses to deeper tissue layers such as muscles and bones.

A visibly noticeable indication of these processes can be non-blanching skin redness, caused by deeper inflammation and capillary damage. If the pressure remains untreated, it can lead to complete necrosis of the affected tissue. Wound infection represents another complication that can occur when pathogens invade the damaged tissue.

Thus, the pathophysiology of pressure ulcers involves a complex interaction of mechanical forces and biochemical processes that ultimately lead to tissue damage. A deep understanding of these processes is crucial to effectively develop preventive and therapeutic measures.

2.3 Risk Factors for the Development of Pressure Ulcers

Decubitus is not caused solely by physical influences such as pressure and shear forces. Various risk factors contribute to the development of pressure ulcers and should always be considered in daily care.

A key factor is limited mobility. Immobility, caused by bedridden conditions or reduced mobility, leads to prolonged static pressure on certain body areas. This increases the risk of inadequate blood circulation and consequently tissue breakdown.

Another risk factor is nutritional status. Inadequate intake of proteins, vitamins, and minerals weakens skin and tissue structure and slows down regenerative processes. Malnutrition and dehydration impair the immune system, increase susceptibility to skin damage, and prolong wound healing time.

Increased skin moisture, caused by incontinence or heavy sweating, also raises the risk of developing decubitus. Moist skin is softer and loses its protective function more quickly. As a result, macerations occur, damaging the skin barrier and serving as entry points for germs.

Conditions such as diabetes mellitus or vascular disorders also affect susceptibility to decubitus. Diabetes can lead to micro- and macroangiopathies, impairing blood circulation and hindering wound healing. Arterial or venous circulation disorders worsen the supply of tissues with oxygen and nutrients.

Sensory impairments, as seen in neurological disorders, also play a role. Patients with reduced pain sensitivity often do not notice pressure points in time, increasing the risk of worsening.

Overall, the risk factors for decubitus are diverse and require critical assessment by caregivers. A comprehensive risk analysis enables individualized prevention planning and targeted measures for decubitus prophylaxis.

2.4 Classification of Pressure Ulcer Stages

2.4.1 Stage 1: Reddening of the skin

A non-blanchable redness of the skin characterizes the first stage of a pressure ulcer. This skin change often occurs at prominent bone tips and body areas where the skin is subjected to increased pressure and shear forces, such as the sacrum, heels, or elbows.

The redness of the skin is the earliest sign of tissue damage that can progress to more severe stages of pressure ulcers without prompt and consistent intervention. The difference from a simple skin redness is that when pressure is applied, the red area does not change color, indicating capillary microthrombosis and beginning inflammatory processes.

Without pressure relief, the damage progresses. Therefore, the rapid recognition of this stage is crucial. Caregivers should conduct regular skin inspections on at-risk residents, specifically looking for non-blanchable redness.

The most important measures for treatment at this stage are pressure relief and the elimination of shear forces. Immediate repositioning of the affected body part is necessary to prevent further damage. Adjustments in positioning techniques, such as the use of soft positioning systems, additionally help to reduce pressure.

Also important is the adherence to consistent skin care. Balance between dryness and moisture is necessary to maintain the natural protective function of the skin. The use of pH-neutral soaps and re-lipidizing creams can support the skin barrier.

The stage also requires increased awareness among care staff and those affected. Appropriate training in recognition and prevention is essential. Mobilization and promotion of self-movement of the affected individuals can be used preventatively to enhance circulation and maintain skin health.

Overall, the first stage represents a critical phase that sets the course for the progression of a pressure ulcer. Early intervention can prevent the transition to more severe stages and significantly improve the quality of life for those affected.

2.4.2 Stage 2: Skin Defect

In stage 2 of pressure ulcers, partial tissue loss is characteristic. This affects the epidermis and can extend into the dermis, often visible as a shallow ulcer or blister. The lesion has a pink-red wound bed color and no necrotic tissue.

Blisters may be intact or open, resulting in an open wound. This stage often occurs in areas of the body that are subjected to constant pressure, such as the buttocks or heels.

Diagnosis in this stage requires a thorough inspection and palpation of the affected skin areas. It is important to protect intact blisters; open blisters are treated as wounds. The use of transparent or hydrocolloid dressings is common as they support a moist wound healing process.

The main goal of treatment includes pressure relief and maintaining a moist wound environment. Continuous monitoring for signs of infection is essential. Equally important is nutritional support through a protein- and energy-rich diet to optimize healing.

Regular repositioning and the use of positioning aids that distribute and reduce pressure are crucial preventive measures. Also, pay attention to skin care. Skin cleansing should be done gently, and moisturizers should be used to ensure barrier function.

For caregivers, training in dealing with affected individuals and early recognition of this stage's features is crucial. Have comprehensive knowledge in wound treatment techniques to prevent progression to stage 3.

Close collaboration in an interdisciplinary team is required to ensure a comprehensive care plan that addresses the specific needs of the individual.

2.4.3 Stage 3: Deeper Tissue Damage

In stage 3 of pressure ulcers, there is a complete loss of skin layers down to the subcutaneous tissue. These deep tissue injuries clinically appear as deep, open wounds with or without visible subcutaneous fat tissue. The adjacent zones of tissue damage are often necrotic and covered with yellowish to grayish fibrin. The wound edges are usually not clearly defined.

The exposure of subcutaneous tissue poses an increased risk of infection, which is why wound care is of utmost priority. Ideally, the wound is cleaned with sterile saline solution to remove necrotic material and debris. Modern wound care materials are used, tailored to the moisture balance of the wound, such as alginates or foam dressings. These materials promote autolytic wound cleaning and create an optimal environment for healing.

In addition to wound care, pressure relief is a central measure. The deep damages originate from continued pressure and shear stress. Regular repositioning and the use of specialized positioning aids and pressure-relieving mattresses are therefore essential. The affected skin area must be kept specifically free from pressure.

Multidisciplinary collaboration is crucial at this stage. This includes consulting wound therapists, specialists, and nutritionists. Any possible systemic cause or comorbidity that impairs wound healing should be treated. The nutritional status of the affected individual must be optimized, with an adequate intake of proteins, vitamins, and calories being necessary.

Thorough documentation of the wound condition, interventions, and healing progress is essential to assess treatment success and make adjustments to the care plan. These records must be regularly reviewed and updated.

Dealing with deep pressure ulcers is challenging for caregivers. Adequate education and training in the area of modern wound care and preventive care interventions are necessary to ensure appropriate care. The goal is to enhance the quality of life for those affected through effective measures and to promote the healing of severely damaged tissues.

2.4.4 Stage 4: Deep Tissue Injury

In stage 4 of a decubitus ulcer, there are profound tissue damages that can extend beyond the subcutaneous layers to muscles, bones, and tendons. These deep wounds are serious medical conditions that require intensive attention and treatment.

The wound often has visible signs of tissue loss down to muscles and bones, possibly associated with necrosis. A wound cavity may develop, characterized by the destruction of deep structures. Extensive slough or necrotic tissue is often found in the wound, which must be carefully removed to promote healing.

Thorough wound cleansing through debridement, which can be performed either surgically or enzymatically, is necessary to remove necrotic material. This action reduces the bacterial load and promotes tissue granulation. The use of advanced wound care techniques such as vacuum therapy can help support the healing process and draw the wound edges together.

Pressure relief remains a crucial measure to promote healing and prevent further damage. Advanced positioning systems and pressure-distributing mattresses are essential to reduce the strain on affected tissues. Hospitals and care facilities should ensure that sufficient specialized aids are available.

At this stage, the risk of infection is high, so regular monitoring for signs of infection is imperative. Adequate antibiotic therapy may be necessary if a systemic or local infection is suspected. Targeted antimicrobial treatment should be aligned based on wound culture results.

The patient's nutrition plays a vital role in the healing process. A diet rich in proteins, vitamins, and minerals supports wound healing and the rebuilding of damaged tissues. Collaboration with nutritionists is advisable to develop an individual nutrition plan.

The complex care requires close collaboration with an interdisciplinary team of professionals, including wound management specialists, nutritionists, and physiotherapists. This ensures comprehensive care and a holistic approach to restoring tissue health.

Caregivers must be familiar with the challenges and techniques of wound treatment. Continuing education in modern wound care is essential to know and apply the latest practices. The focus is on improving the affected individual's quality of life and maintaining independence as much as possible. Detailed documentation of care and healing progress is necessary for the continuous evaluation and adjustment of care plans. These records should be regularly reviewed by all persons involved in the care.

7 This is just a preview

The complete book is available at <https://www.altenpflege.de> for registered users. Registration and use are free of charge. Those who do not want to register can transfer the amount mentioned on page 2 via PayPal to buchhaltung@factsoft.de and include both the book title and email address. They will then receive the PDF via email.

In addition to free access to the short textbooks, registering on [altenpflege.de](https://www.altenpflege.de) enables you to apply directly for jobs and also post a free job advertisement yourself. Become part of the [altenpflege.de](https://www.altenpflege.de) community too.