

altenpflege.de Edition

Short Textbooks for Nursing Education - Focus on Elderly Care

Theo Ahrens

Pathology of Diabetes

presented by

YES
ARCH
ITEC
TURE.

Bibliographic Information

Theo Ahrens – Short Biography

Theo Ahrens, born in 1979 near Munich, is a teacher for nursing professions and a dedicated nursing educator. After completing a voluntary social year in a hospital, he trained as a healthcare and nursing professional, gaining valuable experience in the intensive care unit. Driven by his desire to share knowledge, he studied nursing education and began teaching at a vocational school in Munich in 2009. Theo is known for his practical and empathetic teaching style, preparing future nurses for their demanding profession. Privately, he lives with his wife and two children in Berlin and volunteers to improve conditions in nursing education.

This guide was formulated from his existing materials using AI. The translation was done automatically.

Published by altenpflege.de Price free of charge. Copyright 2024 altenpflege.de Suggestions and improvements please to team@altenpflege.de.

1/1-2412181526

preview

Table of Contents

1 Introduction	4
1.1 Objectives of This Short Textbook	4
1.2 Significance of Diabetes in Elderly Care	4
1.3 Role of Caregivers in Managing Diabetes	5
2 Anatomy and Physiology of Sugar Metabolism	6
2.1 Pancreas and Its Functions	6
2.2 Insulin and Glucagon Function in Glucose Metabolism	6
2.3 Blood Sugar Regulation	7
2.4 Difference between Glucose Uptake and Glucose Utilization	8
3 Forms of Diabetes Mellitus	9
3.1 Type 1 Diabetes	9
3.2 Type 2 Diabetes	9
3.3 Special Forms of Diabetes	10
3.4 Comparison of Types of Diabetes	11
4 Causes and Risk Factors of Diabetes	13
4.1 Genetic Factors	13
4.2 Lifestyle Factors	13
4.3 Age-Related Metabolic Changes	14
4.4 Pharmacological Influences	15
5 Symptoms of Diabetes	16
5.1 Common Symptoms of Type 1 and Type 2 Diabetes	16
5.2 Differences in Symptoms Between Type 1 and Type 2	16
5.3 Slow vs. Sudden Development of Symptoms	17
5.4 Recognition of Warning Signs and Emergency Symptoms	18
6 Diagnosis of Diabetes	19
6.1 Anamnesis	19
6.2 Blood Glucose Measurement	19
6.3 Oral Glucose Tolerance Test (OGTT)	20
6.4 Differences in Diagnostics for Type 1 and Type 2 Diabetes	21
7 Complications and Secondary Diseases in Diabetes	23
7.1 Acute Complications	23
7.2 Long-term Consequences of Diabetes	24
7.3 Neuropathies	25
7.4 Diabetic Foot Syndrome	26
7.5 Retinopathy (Eye Damage)	26
7.6 Nephropathy (Kidney Damage)	28
7.7 Vascular Diseases	29
8 Therapy of Diabetes	30
8.1 Pharmacological Therapy in Type 1 Diabetes	30
8.2 Pharmacological Therapy for Type 2 Diabetes	31
8.3 Types of Insulin and Their Action Profiles	32
8.4 Non-Pharmacological Therapy	33
8.5 Importance of Self-Monitoring Blood Glucose	33
8.6 Technological Support	34
9 Care for Diabetes	36
9.1 Observation of Blood Sugar Levels and Symptoms	36
9.2 Assistance with Conducting Blood Sugar Measurements	36
9.3 Assistance with Insulin Administration	37
9.4 Care of the Diabetic Foot	38
9.5 Nutritional Support	39
9.6 Support with Personal Hygiene	40
9.7 Managing Complications	40
10 Emergency Situations in Diabetes	42
10.1 Hypoglycemia	42
10.2 Hyperglycemia Measures and Emergency Plan	42
10.3 Diabetic Coma	43
10.4 Emergency Call and Collaboration with Emergency Services	44
11 Prevention of Diabetes and Secondary Diseases	46
11.1 Primary Prevention	46
11.2 Secondary Prevention	46
11.3 Prevention of Long-term Complications	47
11.4 Prevention of Diabetic Foot Syndrome	48

1 Introduction

1.1 Objectives of This Short Textbook

This concise textbook provides detailed knowledge about diabetes in the context of elderly care. It serves as a guide to link theoretical foundations with practical applications. Caregivers should be enabled to understand the anatomy and physiology of carbohydrate metabolism to derive informed care decisions from it.

Another goal is the nuanced examination of different forms of diabetes and their specific care approaches. The textbook aims to illuminate the causes and risk factors of diabetes and enable targeted implementation of preventive measures.

An understanding of the symptoms and diagnostic procedures of diabetes is developed, to recognize and prevent complications at an early stage. Caregivers should be enabled to ensure effective care and support of patients through knowledge of therapy options and self-monitoring of blood glucose levels.

Managing emergency situations and the management of secondary diseases are also a focus of this book. The aim is to provide caregivers with tools to act competently and confidently in dealing with diabetic patients and to sustainably improve the well-being of those affected.

1.2 Significance of Diabetes in Elderly Care

Diabetes mellitus in geriatric care

Diabetes mellitus is of central importance in geriatric care. This condition often affects older people, so geriatric caregivers regularly encounter this metabolic disorder. Managing diabetes requires specific knowledge and skills, as older people face additional challenges due to age-related changes in the body.

The care of people with diabetes in geriatric care includes monitoring blood sugar levels, assisting with medication intake and insulin administration, and implementing measures to prevent secondary diseases and complications. Caregivers play a crucial role in implementing individualized care plans tailored to the specific needs of the residents.

Since diabetes in the elderly often occurs alongside other chronic conditions such as hypertension and cardiovascular diseases, holistic care is necessary. Caregivers must be able to recognize interactions between diseases and the medications used and act accordingly. Good collaboration with doctors and other healthcare professionals is essential to achieve optimal care outcomes.

The importance of diabetes in geriatric care is also evident in the necessity of preventing acute complications such as hypoglycemia and hyperglycemia, which can have serious consequences in the elderly. Caregivers should be able to recognize warning signs early and respond appropriately to ensure the safety of residents.

Overall, diabetes as one of the most common conditions in old age will continue to shape care practice. It requires a comprehensive understanding of the disease and its impact on the daily routine of geriatric care to sustainably improve the quality of life for those affected.

1.3 Role of Caregivers in Managing Diabetes

Caregivers undertake versatile tasks in managing diabetes. Their primary role is to continuously monitor the blood sugar levels of residents. Regular measurements help detect deviations early and initiate necessary actions to prevent dangerous complications.

Another central aspect is the support in administering insulin and other antidiabetic medications. Caregivers ensure that medications are correctly dosed and administered timely according to medical orders. This requires special care and knowledge about the medication's effects and potential side effects.

Caregivers also serve as intermediaries between residents, doctors, and family members. They are often the first point of contact for questions or concerns about the illness and must possess well-founded knowledge to provide comprehensive advice and information. It is important to communicate clearly and understandably to avoid misunderstandings.

Nutritional counseling and motivation for lifestyle changes also belong to their tasks. Caregivers assist residents in creating an individual nutritional plan and promote physical activity within the limits of the respective health condition. The aim is to enable residents to independently contribute to stabilizing their blood sugar levels.

Additionally, caregivers are responsible for the observation and care of potential diabetes complications, such as diabetic foot syndrome. Regular foot checks and conscientious documentation are crucial to notice tissue damage early and initiate measures to improve healing.

Finally, caregivers are qualified for managing emergency situations. They recognize and intervene in cases of hypoglycemia or hyperglycemia and activate emergency services if necessary. Through targeted training and continuous education, they keep their knowledge up-to-date to act safely and competently in critical situations.

Overall, caregivers significantly contribute to the quality of life, safety, and well-being of residents with diabetes. Their work requires a high level of expertise, precision, and empathy.

2 Anatomy and Physiology of Sugar Metabolism

2.1 Pancreas and Its Functions

The pancreas, also known as the pancreas, is a central organ for sugar metabolism. It is located in the upper abdomen, directly behind the stomach, and has both endocrine and exocrine functions.

The endocrine part of the pancreas is responsible for hormone production. Here, the islets of Langerhans are located, where specific cell types produce insulin and glucagon. Beta cells in the islets of Langerhans release insulin, a hormone that lowers blood sugar levels. Insulin promotes the uptake of glucose into body cells, thus supporting energy metabolism. Alpha cells, on the other hand, produce glucagon, which raises blood sugar levels. Glucagon stimulates the liver to release stored glucose into the bloodstream. These two hormones act as antagonists, working together to regulate blood sugar levels.

The exocrine part of the pancreas produces digestive enzymes that are released into the small intestine. These enzymes are essential for the digestion of fats, proteins, and carbohydrates, but are not directly related to blood sugar regulation.

Functional disorders in the endocrine area of the pancreas, whether due to autoimmune processes, inflammation, or other damage, can lead to the development of diabetes mellitus. In type 1 diabetes, the body's own immune system destroys the insulin-producing beta cells, while in type 2 diabetes, insulin resistance of the body cells is often at the forefront. The result in both cases is elevated blood sugar levels, which can lead to various health problems in the long run.

The function of the pancreas is therefore of great importance not only for sugar metabolism but also for the overall well-being of the body. Understanding the processes that occur in this organ is essential to better comprehend the development and treatment of diabetes mellitus.

2.2 Insulin and Glucagon Function in Glucose Metabolism

Insulin and glucagon are central hormones in glucose metabolism that work in a finely tuned balance to ensure stable blood sugar levels. Insulin is released after food intake when blood sugar levels rise. It promotes the uptake of glucose into muscle and fat cells, allowing its use for energy production or its storage as glycogen in the liver. Additionally, insulin prevents the new production of glucose in the liver, which also contributes to lowering blood sugar.

Glucagon, on the other hand, is primarily released during periods of fasting or when blood sugar levels are low. It activates the conversion of glycogen into glucose in the liver to release it into the blood and increase blood sugar levels. Additionally, glucagon promotes gluconeogenesis, a process in which new glucose is formed in the liver from non-carbohydrate precursors.

The interaction of insulin and glucagon can be compared to a thermostat system that maintains temperature consistency in a room. When blood sugar rises, insulin "switches on" and lowers the level. If blood sugar drops too much, glucagon kicks in and raises it. This dynamic regulation allows for a continuous and reliable flow of energy to the body's cells.

Disruptions in the function or regulation of insulin and glucagon can lead to significant health problems. In type 1 diabetes, insulin is completely absent, which, if untreated, results in highly elevated blood sugar levels and an energy deficit in cells. In type 2 diabetes, the action of insulin on the cells is reduced (insulin resistance), while insulin production may initially persist. In both cases, the precise balance between insulin and glucagon is disturbed, necessitating modern therapeutic measures. A deep understanding of the function of these hormones allows for the identification of appropriate treatment and care approaches, improving the quality of life for affected individuals.

2.3 Blood Sugar Regulation

Blood Sugar Regulation

Blood sugar regulation is a complex interplay of numerous processes that keep the blood sugar level stable within narrow limits. Normally, fasting blood sugar levels for adults range between 70 and 99 mg/dl. After food intake, it may rise temporarily but should quickly return to normal levels through the body's own mechanisms.

Central to the regulation of blood sugar levels are the hormones insulin and glucagon, produced in the pancreas. As previously described, insulin promotes the uptake of glucose into the cells when blood sugar levels rise after food intake, thereby reducing blood sugar. At the same time, it blocks the new production and release of glucose in the liver. In contrast, glucagon is released when blood sugar levels drop to stimulate the formation and release of glucose from the liver's glycogen stores.

In addition to these main players, other hormones also contribute to blood sugar regulation. Adrenaline and cortisol, stress hormones, can increase blood sugar levels by supporting glucose release from the liver. Growth hormone, on the other hand, acts by modifying the cells' sensitivity to insulin.

A functioning regulation system ensures that glucose supply optimally meets the body's energy needs. In cases of dysfunction, as seen in diabetes mellitus, this balance is disrupted. This leads to blood sugar fluctuations that, if untreated, can cause acute complications in the short term and long-term damage to organs and tissues. Understanding the mechanisms of blood sugar regulation is therefore fundamental for recognizing and treating such disorders and for supporting affected patients in everyday life.

2.4 Difference between Glucose Uptake and Glucose Utilization

Glucose uptake and glucose utilization are two central processes in carbohydrate metabolism that work seamlessly together to provide energy for the body and maintain balanced blood sugar levels.

Glucose uptake refers to the transport of glucose from the blood into the body's cells. This predominantly occurs under the influence of insulin, a hormone from the pancreas. Insulin opens "transport gates" (glucose transporters), allowing glucose to enter muscle and fat cells, where it is either used for energy production or stored as glycogen. The insulin-dependent glucose transport is crucial for lowering blood sugar levels after meals.

In contrast, glucose utilization describes the process by which cells use the absorbed glucose for their energy needs. This process, also known as cellular respiration, occurs in various steps within the cells: First, glucose is broken down into pyruvate through glycolysis. In a subsequent step, the citric acid cycle, the complete oxidative breakdown to carbon dioxide and water occurs, releasing stored energy in the form of ATP (adenosine triphosphate).

The two processes are closely linked. Efficient glucose uptake into the cells is a prerequisite for effective glucose utilization and thus for maintaining a stable energy metabolism. If glucose uptake is disrupted, for example, by insulin resistance or insulin deficiency, glucose remains in the blood, leading to elevated blood sugar levels, as observed in diabetes.

An example illustrates this difference: After a meal, blood sugar levels rise. Insulin is released to stimulate glucose uptake. If this works, glucose quickly enters the cells and is utilized, normalizing blood sugar levels. In case of a disruption of these processes, such as impaired insulin action, blood sugar levels remain high, as glucose uptake does not proceed as expected and utilization cannot occur.

A deeper understanding of these processes is crucial to plan and effectively implement preventive and therapeutic measures in diabetes.

3 Forms of Diabetes Mellitus

3.1 Type 1 Diabetes

Type 1 diabetes is an autoimmune disease where the body's immune system attacks the insulin-producing beta cells in the pancreas. This destruction leads to an absolute insulin deficiency. Since insulin is necessary to transport glucose from the blood into the body's cells, a persistently high blood sugar level results.

Causes of this autoimmune process are largely genetically determined, although environmental factors can also be involved. A precise cause is often difficult to determine. Affected individuals often have no direct family history, indicating complex genetic relationships.

Type 1 diabetes typically presents acutely. Typical symptoms include excessive thirst, frequent urination, fatigue, and weight loss. These symptoms develop quickly, usually within a few weeks. The rapid destruction of beta cells necessitates immediate insulin therapy.

The course of type 1 diabetes is characterized by the fact that affected individuals are dependent on insulin administration for life. It requires a precisely tailored insulin therapy, adjusted to the individual's metabolic condition. A challenge for caregivers is to support patients in self-monitoring their blood sugar levels and adjusting the insulin dosage.

Poor blood sugar control can lead to acute problems such as ketoacidosis and, in the long term, complications such as damage to the eyes, kidneys, and nerves. Therefore, close monitoring and adjustment of treatment are crucial. Caregivers play a significant role in educating those affected and supporting them in their daily lives.

3.2 Type 2 Diabetes

Type 2 Diabetes is a metabolic disorder characterized by insulin resistance and consequently a relative insulin deficiency. Unlike Type 1, the body does produce insulin in Type 2, but the body's cells are less sensitive to it. This results in glucose remaining in the blood and the blood sugar level rising.

Common causes for the development of Type 2 diabetes include genetic predispositions as well as lifestyle factors such as lack of exercise, unhealthy diet, and overweight. Particularly, visceral fat surrounding the organs in the abdominal cavity plays a significant role in the development of insulin resistance. Age-related metabolic changes may also increase the risk, especially among the older population, which is the focus in geriatric care.

The symptoms of Type 2 diabetes often develop gradually and can remain unnoticed over many years. Indicators include increased thirst, frequent urination, fatigue, and nonspecific symptoms like vision problems or slow-healing wounds. Many affected individuals only become aware of the disease at an advanced stage or when complications arise.

The progression of Type 2 diabetes is chronic and progressive. Without appropriate treatment, serious complications can occur, including cardiovascular diseases, kidney damage, and vision disorders. Insulin therapy is often not required initially. Treatment typically begins with lifestyle changes — nutrition and exercise — and, if necessary, the intake of oral antidiabetic medications. Caregivers play a crucial role in supporting patients, particularly through education on lifestyle changes, medication monitoring, and promoting adherence to therapy. The long-term goal is to control blood sugar levels and maintain the quality of life for those affected.

3.3 Special Forms of Diabetes

Gestational Diabetes

Gestational diabetes occurs during pregnancy and is one of the most common metabolic disorders at this time. This form of diabetes is characterized by insulin resistance, which is triggered by hormonal changes. Since the body needs to produce more insulin to regulate the increased blood sugar levels, inadequate insulin production can lead to elevated glucose levels.

Typical risk factors include being overweight, older maternal age, or a family history of type 2 diabetes. Gestational diabetes is usually diagnosed through an oral glucose tolerance test conducted between the 24th and 28th week of pregnancy.

Careful monitoring and, if necessary, dietary adjustments are crucial to regulate blood sugar levels. In severe cases, insulin therapy may be required. After birth, blood sugar levels usually normalize, but the risk of developing type 2 diabetes later remains increased.

MODY - Maturity Onset Diabetes of the Young

MODY is a rare, genetically determined form of diabetes where mutations in individual genes lead to disruptions in insulin production. Unlike type 1 or type 2 diabetes, MODY typically manifests before the age of 25 and is not directly associated with autoimmune processes or insulin resistance.

There are various MODY subtypes, differing in the cause of the genetic mutation. A key feature is that affected individuals often appear across several generations in a family, making genetic counseling advisable.

Treatment depends on the specific type of MODY and can range from dietary measures and oral antidiabetics to insulin therapy. Correct diagnosis and differentiation from type 1 or type 2 diabetes are crucial for the appropriate therapeutic strategy.

7 This is just a preview

The complete book is available at <https://www.altenpflege.de> for registered users. Registration and use are free of charge. Those who do not want to register can transfer the amount mentioned on page 2 via PayPal to buchhaltung@factsoft.de and include both the book title and email address. They will then receive the PDF via email.

In addition to free access to the short textbooks, registering on [altenpflege.de](https://www.altenpflege.de) enables you to apply directly for jobs and also post a free job advertisement yourself. Become part of the [altenpflege.de](https://www.altenpflege.de) community too.